

Wetting Transitions of Methanol on the n-Alkanes

D. Ross and M. Moldover

National Institute of Standards and Technology

Physical and Chemical Properties Division

100 Bureau Drive, Stop 8380

Gaithersburg, MD 20899-8380 U.S.A.

D. Bonn and J. Meunier

Laboratoire de Physique Statistique

Ecole Normale Supérieure

24 rue Lhomond

75231 Paris cedex 05, France

Results are presented of the wetting properties of methanol (CH_3OH) at the liquid-vapor interface of different n-alkanes ($\text{C}_n\text{H}_{2n-2}$, denoted as C_n) ranging from hexane (C_6) to undecane (C_{11}) as well as alkane mixtures with non-integer effective carbon numbers ranging from $n = 8$ to 10. Measurement of the contact angle shows that, as the bulk critical point is approached, wetting transitions occur for long alkanes ($n > 8$), whereas drying is found for short alkanes ($n < 8$). Measurements of the wetting layer thickness and precise contact angle measurements close to the wetting temperature, T_w , reveal a change in the order of the transition. First-order (discontinuous) wetting occurs for T_w far from the bulk critical point T_c . Critical (continuous) wetting is observed for T_w close to T_c . The observed critical wetting transition has all the features of the long-sought short-range critical wetting transition. The results for the critical wetting transition are discussed in detail.